Demo Our Full Clinical Procedure Manual



Tinetti Gait and Balance Test

Balance

Instructions to Examiner

- · Provide overview of what the assessment will consist of and tell them you'll talk them through it
- Begin with patient seated in hard, armless, stably positioned chair
- For rising and sitting, ask patient to attempt without using their arms/hands
- Examiner stands near patient, few feet in front and slightly to the side; spot for safety at all times
- For the nudge test, the patient should stand with feet as close together as possible; examiner presses on sternum with palm of hand (3 trials)

Domain	Evaluation Characteristics		Description of Deficit
Sitting balance	Leans or slides in chair Steady, safe	= 0 = 1	
Rises from chair	Unable to without help Able, uses arms to help Able without use of arms	= 0 = 1 = 2	
Attempts to rise	Unable to without help Able, requires > 1 attempt Able to rise, 1 attempt	= 0 = 1 = 2	
Immediate standing balance (first 5 seconds)	Unable to without help Able, requires > 1 attempt Able to rise, 1 attempt	= 0 = 1 = 2	
Standing balance	Unsteady Steady but wide stance (>4 inches) or uses support	= 0 = 1	
	Narrow stance without support	= 2	
Nudge	Begins to fall Staggers, but catches self Steady	= 0 = 1 = 2	
Nudge, eyes closed	Unsteady Steady	= 0 = 1	
Turning 360 degrees ("make a complete circle")	Discontinuous steps Continuous	= 0 = 1	
	Unsteady (grabs, staggers) Steady	= 0 = 1	
Sitting down	Unsafe (misjudged distance, falls	= 0	
	into chair) Uses arms or not a smooth motion Safe, smooth motion	= 1 = 2	

Demo Our Full Clinical Procedure Manual



Gait

Definitions of terms used in gait assessment tool

Tinetti Gait and Balance Test

- "Aid" is rated positive if patient is using his/her walking aid or if patient grabs at rails or furniture
- "Stance foot" is the foot that remains on the ground during gait cycle

Instructions to Examiner

- Examiner moves along with patient; spot for safety at all times
- Patient walks across room (using walking aid if customary for them) for approximately 25 feet each way
- · Provide overview of what assessment will consist of and tell patient you'll talk them through it
- On way out ask patient to walk at usual pace; on way back ask patient to walk at rapid but safe pace
- · Instruct patient about the starting cue (go), and that they should stop if they feel unsafe

Domain	Evaluation Characteristics		Description of Deficit
Indication of gait (immediately after "'go")	Any hesitancy or multiple attempts No hesitancy	= 0 = 1	
Step length and height	Right foot swing Does not pass L stance foot Steps past L foot Does not clear floor Clears floor Left foot swing Does not pass R stance foot Steps past R foot Does not clear floor Clears floor	= 0 = 1 = 0 = 1 = 0 = 1 = 0 = 1	
Step symmetry	Right and left step length not equal Right and left step length equal	= 0 = 1	
Step continuity	Stopping or discontinuity between steps Steps continuous	= 0 = 1	
Path	Marked deviation Mild/moderate deviation or uses aid Straight without aid	= 0 = 1 = 2	
Trunk	Marked sway or uses aid No sway but flexed knees or back or spread arms wide No sway, flexion, widened arms, or aid	= 0 = 1 = 2	
Walking stance	Heels apart Heels almost touching while walking	= 0 = 1	

Demo Our Full Clinical Procedure Manual



Tinetti Gait and Balance Test

Gait score

Balance score (prior page)

Total score

/12
/16
/28

Clinical procedures optimized for tablets and phones, including:

- Medication Reconciliation in the Home
- Transcutaneous Electrical Nerve Stimulation (TENS) Unit
 - Pain Assessment
 - · Neurological Assessment
 - 100+ more

Easy, fast and more reliable than YouTube or Google



Get a demo today www.medbridge.com

